

SUTAB Instructions for Colonoscopy

Date of procedure: add date Arrive at: add timePlease select

Location of Department:

Ochsner Medical Center Please select **Please select** Please select

As soon as possible:

• Pick up your prep from the pharmacy

On the day before your procedure...

DO NOT:

- Do not EAT solid food, drink milk or anything colored red.
- Do not drink alcohol.
- Do not take other laxatives while taking SUTAB.
- Do not take oral medications within 1 hour of starting each dose of SUTAB.
- No gum chewing or candy morning of procedure
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.

DO:

Drink <u>clear liquids ONLY</u> - Drink at least 6 to 8 glasses of clear liquids from time you wake up until you begin your prep to avoid dehydration.

Liquids That Are **OK** to Drink:

- Water
- Sports drinks (Gatorade, Power-Aid)
- Coffee or tea (no cream or nondairy creamer)
- Clear juices without pulp (apple, white grape)
- Gelatin desserts (no fruit or toppings)
- Clear soda (sprite, coke, ginger ale)
- Chicken broth (until 12 midnight the night before procedure)

***(Please disregard the insert instructions from pharmacy and follow these instructions).

Note:

- SUTAB is an osmotic laxative indicated for cleansing of the colon in preparation for colonoscopy in adults.
- Medication taken by mouth may not be absorbed properly when taken within 1 hour before the start of each dose of SUTAB.

<u>IMPORTANT:</u> If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause, or slow the rate of drinking the additional water until your symptoms decrease.

How to take prep:

SUTAB is a (2-day) prep. A total of 24 tablets are required for complete preparation for colonoscopy.

DOSE 1—Day Before Colonoscopy add date at add time.

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow 1 tablet every 1 to 2 minutes. You should finish the 12 tablets and the entire 16 ounces of water within 20 minutes.

STEP 3 After 1 hour of completing step 2, drink at least 32 ounces of additional water/clear liquids until bedtime, but more is better.

DOSE 2—Day of the Colonoscopy add date at Please select.

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow 1 tablet every 1 to 2 minutes. You should finish the 12 tablets and the entire 16 ounces of water within 20 minutes.

STEP 3 After 1 hour of completing step 2, drink at least 32 ounces of additional water/clear liquids until 4 hours before your colonoscopy then nothing else by mouth or as directed by the scheduling nurse add time.

For more information about your procedure, please watch this informational video.

Options for viewing:

Using a keyboard: press and hold the control tab
(Ctrl) and left mouse click to follow link
Colonoscopy Video – YouTube

OR

- Type link address into your web browser's address bar: https://www.youtube.com/watch?v=XZdo-
- Using a mobile phone: tap on web address/link.

Comments:	

LP1xDQ