

Sugar & Breathing & Sepsis...Oh my!

The Neonatal Transition to the Outside World

Amanda England, MD, MPH

July 19, 2019

Financial Disclosures

- None

Outline

- Hypoglycemia
 - Normal physiology
 - Screening
 - Treatment
- Respiratory Distress
 - Transition from fetal life
 - Common causes of neonatal respiratory distress
- Sepsis- AAP December 2018 Policy

Physiology

- Glucose supplied via mother
 - Glucose is the main energy supply of the fetus
- Fetus stores glucose as glycogen
 - Glycogen stored in liver, heart, lung, and skeletal muscle
- Cut umbilical cord triggers glycogenolysis, gluconeogenesis, and utilization of exogenous sources via feeding
- Glucose nadir usually between 1-2 hours after delivery

Hypoglycemia Etiologies

- Increased utilization of glucose
 - Hyperinsulinism
 - Without hyperinsulinism
- Decreased glucose production

Hyperinsulinism

- Infants of diabetic mothers
- LGA
- Beckwith-Wiedemann syndrome
- Insulin producing tumors
- Maternal drugs

Maternal Medications

Medication	Used for Treatment of
β-sympathomimetics: Terbutaline	Tocolytic therapy
Sulfonylureas: Glyburide, Glipizide	Type 2 diabetes
β-blockers: Labetalol, Propranolol, Metoprolol	Hypertension, migraine headaches, thyrotoxicosis
Thiazide diuretics: Chlorothiazide, Hydrochlorothiazide	Hypertension, edema
Tricyclic antidepressants: Amitriptyline, Nortriptyline, Desipramine	Depression

Without Hyperinsulinism

- Perinatal stress
 - Sepsis
 - Shock
 - Asphyxia
 - **Hypothermia**
- Polycythemia
- Endocrine deficiencies
- Metabolic disorders

Decreased Glucose Production/Stores

- Prematurity
- IUGR/SGA
- Inadequate caloric intake
- Delayed onset of feeding

AAP Guidelines

- No screening is necessary for healthy, term neonates in the setting of a normal pregnancy
- Risk Factors:
 - SGA
 - IDM
 - LGA
 - Preterm
- Infant with clinical signs of hypoglycemia

AAP Guidelines- Clinical Signs

- Jitteriness
- Cyanosis
- Seizures
- Apneic episodes
- Tachypnea
- Weak or high-pitched cry
- Floppiness or lethargy
- Poor feeding
- Eye rolling

AAP Guidelines- When to screen?

- ASAP- minutes, not hours
 - At risk infants should be fed within 1 hour of life
 - Screen 30 minutes after first feeding
- Screening for 12 hours:
 - IDM
 - LGA
- Screening for 24 hours:
 - SGA
 - Late preterm infants

Screening and Management of Postnatal Glucose Homeostasis in Late Preterm and Term SGA, IDM/LGA Infants

[(LPT) Infants 34 – 36^{6/7} weeks and SGA (screen 0-24 hrs); IDM and LGA ≥ 34 weeks (screen 0-12 hrs)]

Symptomatic and <40 mg/dL $\longrightarrow$ IV glucose

ASYMPTOMATIC

Birth to 4 hours of age

INITIAL FEED WITHIN 1 hour
Screen glucose 30 minutes after 1st feed

Initial screen <25 mg/dL

Feed and check in 1 hour

<25 mg/dL

↓
IV glucose*

25–40 mg/dL

↓
Refeed/IV glucose*
as needed

4 to 24 hours of age

Continue feeds q 2-3 hours
Screen glucose prior to each feed

Screen <35 mg/dL

Feed and check in 1 hour

<35 mg/dL

↓
IV glucose*

35 – 45 mg/dL

↓
Refeed/IV glucose*
as needed


Target glucose screen ≥ 45 mg/dL prior to routine feeds


* Glucose dose = 200 mg/kg (dextrose 10% at 2 mL/kg) and/or IV infusion at 5–8 mg/kg per min (80–100 mL/kg per d). Achieve plasma glucose level of 40-50 mg/dL.

Symptoms of hypoglycemia include: Irritability, tremors, jitteriness, exaggerated Moro reflex, high-pitched cry, seizures, lethargy, floppiness, cyanosis, apnea, poor feeding.

Hypoglycemia Treatment







Transition from Fetal Life


- Replacement of alveolar fluid with air
- Onset of regular breathing
- Increase in pulmonary blood flow
 - Increase in systemic vascular resistance
 - Decrease in pulmonary vascular resistance

Risk Factors for Difficulty in Transition

- Maternal Factors:
 - AMA
 - Maternal DM or HTN
 - Substance abuse
 - Previous history of stillbirth or early neonatal loss
- Fetal Factors:
 - Prematurity
 - Multiples
 - Congenital anomalies
- Antepartum Conditions:
 - Placental anomalies
 - Oligohydramnios/Polyhydramnios
- Delivery Complications:
 - Transverse or breech
 - Chorioamnionitis
 - Meconium-stained fluid
 - Abnormal fetal heart tracing
 - Maternal narcotics within 4 hours
 - Instrument-assisted delivery

Transient Tachypnea of Newborn

- Delayed resorption of fetal lung fluid
 - Increase risk in infants delivered via elective Cesarean section
- Pulmonary immaturity
- CXR:
 - Hyperinflation
 - Perihilar fullness/streaking
 - Fluid in the minor fissure
 - Occasional effusion



Transient Tachypnea of Newborn

- Clinical Presentation:
 - **Tachypnea**
 - Increased work of breathing- grunting, retractions, nasal flaring
 - Barrel chest
 - Auscultation- good air entry +/- crackles
 - Onset within the first 2 hours of life
- Symptoms can last 12-72 hours

Transient Tachypnea of Newborn

- Management:
 - Supportive
 - Oxygen
 - Provide lung recruitment- High flow nasal cannula, CPAP
 - Gavage feedings with significant tachypnea
 - No role for diuretics

Respiratory Distress Syndrome


- Surfactant deficiency
- Surfactant produced by type II pneumocytes
 - These cells differentiate around weeks 24-28
- Maintains alveolar stability

Respiratory Distress Syndrome

- Risk Factors:
 - **Prematurity**
 - Male
 - Maternal diabetes
 - C-section without labor
 - Perinatal asphyxia
 - Chorioamnionitis


Respiratory Distress Syndrome

- Clinical Presentation:
 - Respiratory distress at birth
 - Worsens before it improves
 - CXR- ground glass appearance with air bronchograms
- Management:
 - Antenatal steroids
 - PEEP
 - Artificial surfactant administration



Persistent Pulmonary Hypertension

- Pulmonary vascular resistance remains elevated
- Pathophysiology:
 - Underdevelopment
 - Maladaptation
 - Maldevelopment



Persistent Pulmonary Hypertension

- Risk Factors
 - Meconium aspiration
 - Congenital anomalies
 - Pneumonia
 - Sepsis
 - Asphyxia/Perinatal hypoxia
 - Polycythemia
 - Congenital heart disease
 - Maternal medications

Persistent Pulmonary Hypertension


- Clinical Presentation:
 - Respiratory distress with hypoxia
 - Differential saturations
 - CXR- “black lung”
 - Murmur
- Echocardiogram



Persistent Pulmonary Hypertension

- Management:
 - Oxygen!
 - This may require intubation
 - Correct acidosis
 - Hyperventilation and alkalinization no longer recommended
 - Hemodynamic Support
 - iNO
 - Transfer to the NICU/ECMO Center
 - $OI = (FiO_2 * MAP) / PaO_2$





Early Onset Sepsis

- By the numbers
 - Incidence 0.5 in 1000 births
 - Late preterm incidence 1 in 1000 births
 - Culture confirmed meningitis 0.01-0.02 cases per 1000 births
 - Mortality 2-3%
- Usually begins in utero

Early Onset Sepsis

- Risk Factors:
 - < 37 weeks gestation
 - ROM > 18 hours
 - Maternal infection
 - Maternal GBS status
 - Appropriate intrapartum antibiotic administration

Early Onset Sepsis

- Risk stratification:
 - Categorical Risk Factor Assessment
 - Risk Assessment Based on Newborn Condition
 - Multivariate Risk Assessment

Neonatal Early-Onset Sepsis Calculator

Kaiser Permanente
Research

Home Classification References

Probability of Neonatal Early-Onset Sepsis Based on Maternal Risk Factors and the Infant's Clinical Presentation

The tool below is intended for the use of clinicians trained and experienced in the care of newborn infants. Using this tool, the risk of early-onset sepsis can be calculated in an infant born ≥ 34 weeks gestation. The interactive calculator produces the probability of early onset sepsis per 1000 babies by entering values for the specified maternal risk factors along with the infant's clinical presentation.



Please enter details below.

Predictor

Scenario

Calculate »

Clear

Please enter details below.

Predictor	Scenario
Incidence of Early-Onset Sepsis ?	<input type="text"/>
Gestational age ?	<input type="text"/> weeks <input type="text"/> days
Highest maternal antepartum temperature ?	<input type="text"/> Fahrenheit
ROM (Hours) ?	<input type="text"/>
Maternal GBS status ?	☐ Negative ☐ Positive ☐ Unknown
Type of intrapartum antibiotics ?	☐ Broad spectrum antibiotics > 4 hrs prior to birth ☐ Broad spectrum antibiotics 2-3.9 hrs prior to birth ☐ GBS specific antibiotics > 2 hrs prior to birth ☐ No antibiotics or any antibiotics < 2 hrs prior to birth

Calculate » Clear

Risk per 1000/births

EOS Risk @ Birth

EOS Risk after Clinical Exam	Risk per 1000/births	Clinical Recommendation	Vitals
Well Appearing			
Equivocal			
Clinical Illness			

Classification of Infant's Clinical Presentation [Clinical Illness](#) [Equivocal](#) [Well Appearing](#)

Please enter details below.

Predictor	Scenario
Incidence of Early-Onset Sepsis ?	0.5/1000 live births (CDC national) ▼
Gestational age ?	<div>36 weeks</div> <div>2 days</div>
Highest maternal antepartum temperature ?	<div>98.9 Fahrenheit ▼</div>
ROM (Hours) ?	6
Maternal GBS status ?	☒ Negative ☐ Positive ☐ Unknown
Type of intrapartum antibiotics ?	☐ Broad spectrum antibiotics > 4 hrs prior to birth ☐ Broad spectrum antibiotics 2-3.9 hrs prior to birth ☐ GBS specific antibiotics > 2 hrs prior to birth ☒ No antibiotics or any antibiotics < 2 hrs prior to birth

Calculate »

Clear

Risk per 1000/births			
EOS Risk @ Birth	0.37		
EOS Risk after Clinical Exam	Risk per 1000/births	Clinical Recommendation	Vitals
Well Appearing	0.15	No culture, no antibiotics	Routine Vitals
Equivocal	1.82	Blood culture	Vitals every 4 hours for 24 hours
Clinical Illness	7.68	Empiric antibiotics	Vitals per NICU

Classification of Infant's Clinical Presentation	Clinical Illness	Equivocal	Well Appearing
<p>1. Well Appearing</p> <p>2. Clinical Illness</p> <p>3. Equivocal</p>	<p>1. Well Appearing</p> <p>2. Clinical Illness</p> <p>3. Equivocal</p>	<p>1. Well Appearing</p> <p>2. Clinical Illness</p> <p>3. Equivocal</p>	<p>1. Well Appearing</p> <p>2. Clinical Illness</p> <p>3. Equivocal</p>

Please enter details below.

Predictor	Scenario
Incidence of Early-Onset Sepsis ?	0.5/1000 live births (CDC nationa ▾)
Gestational age ?	<div>36 weeks</div> <div>2 days</div>
Highest maternal antepartum temperature ?	<div>98.9 Fahrenheit ▾</div>
ROM (Hours) ?	<div>6</div>
Maternal GBS status ?	<div>☒ Negative</div> <div>☐ Positive</div> <div>☐ Unknown</div>
Type of intrapartum antibiotics ?	<div>☐ Broad spectrum antibiotics > 4 hrs prior to birth</div> <div>☐ Broad spectrum antibiotics 2-3.9 hrs prior to birth</div> <div>☐ GBS specific antibiotics > 2 hrs prior to birth</div> <div>☒ No antibiotics or any antibiotics < 2 hrs prior to birth</div>

Calculate »

Clear

Risk per 1000/births

EOS Risk @ Birth

0.37

EOS Risk after Clinical Exam	Risk per 1000/births	Clinical Recommendation	Vitals
Well Appearing	0.15	No culture, no antibiotics	Routine Vitals
Equivocal	1.82	Blood culture	Vitals every 4 hours for 24 hours
Clinical Illness	7.68	Empiric antibiotics	Vitals per NICU

Classification of Infant's Clinical Presentation Clinical Illness Equivocal Well Appearing

Early Onset Sepsis


- Clinical Manifestations:
 - Fetal/delivery room distress
 - Temperature instability
 - Respiratory distress
 - Cardiocirculatory symptoms
 - Neurologic symptoms

Early Onset Sepsis

- Laboratory Tests:
 - Blood culture
 - Need 1 mL of blood minimum
 - Lumbar puncture
 - CBC
 - Inflammatory markers

Early Onset Sepsis

- Most Common Pathogens:
 - Group B Streptococcus
 - Escherichia coli
 - Gram positive organisms
 - Listeria monocytogenes
 - Staphylococcus aureus



Early Onset Sepsis

- Treatment:
 - Ampicillin and gentamicin
- Culture positive:
 - Narrow treatment after ID
 - Serial repeat blood cultures
- Culture negative:
 - Discontinue after 36-48 hours of therapy

Thank You!

Amanda.England@Ochsner.org