




Neurocognitive Neuroanatomy

Focus on Fasciculus Network
Phenomenology

May 15th, 2024

James Rini MD. MPH.
Assistant Professor
Ochsner Neurocognitive
Disorders Program

Presenter Disclosure

Faculty: James Fraser Rini


I have no current or past relationships with commercial entities.

Potential for conflict(s) of interest: None

Commercial Support Disclosure

To the presenter knowledge, this program has received no financial or in-kind support from any commercial or other organization.

Potential for conflict(s) of interest: None



strategies for AD, by allowing for selection of patients on the basis of biological as well as clinical and epidemiological factors.

A Rose By Any Other Name

Linguistic relativity: Language Impacts How We See


Perceptual Knowledge is **strongly** influenced by learnt distinctions

Sapir-Whorf hypothesis c.1929


- Structure of a language determines a native speaker's perception and categorization of experience.

Himba tribe of northern Namibia

- *Serandu*: red, orange and pink
- *Zoozu*: black and dark shades of red, blue, green and purple
- *Vapa*: white and some yellow
- *Borou*: greens and blues (lacking the blue-green distinction)
- *Dumbu*: other greens, red and brown



Lesson: *You don't see what you haven't been taught to distinguish*



**WHY DOES
IT MATTER?**


Its not the size that matters

Size of CNS generally correlates with general intelligence

BUT relationship is neither robust nor mechanistically understood.

- Several large mammals have substantially larger brains than humans
- *So, did Neanderthal*

Structural and functional differences appear to be the root of functional connectivity and higher levels of consciousness




Connectome

Cognition and behavior are a function dynamic network connectivity i.e. connectome

Human connectome appears to be unique

- Many species-specific differences in the organization
- Short-range projections in association areas are less myelinated
- Long-range projections in prefrontal/temporal are more numerous

Understanding how these multilayered bidirectional tract/bundles are involved in processing and modulating decisions and behaviors is crucial for understanding cognition and higher levels of consciousness



The First Fasciculi



400 million years ago vertebrates evolved early axon scaffolding (i.e. bundles and myelin)

First and most conserved tract is **ventral longitudinal tract (VLT)**


- medial longitudinal fascicle (MLF)
- postoptic commissure (TPOC)

Fasciculi structures evolved for coordinate efficient and effective motor responses to cephalic sensory stimuli

- Vestibulo-ocular (VOR) reflex
- Optokinetic (OKR) reflex
- Phototaxis
- Jaw movement



400 million years later
...things are a complicated



- | | |
|--|--|
| • Cuneate Fasciculus | • Tract |
| • Gracile Fasciculus | • Anterolateral System |
| • Spinothalamic Tract | • Dorsal Premammillary Nucleus |
| • Spinocerebellar Tract | • Lateral Olfactory Tract |
| • Dorsal Spinocerebellar Tract | • Superior Longitudinal Fasciculus (SLF) |
| • Ventral Spinocerebellar Tract | • Arcuate Fasciculus |
| • Dorsal Column-Medial Lemniscus Pathway | • Uncinate Fasciculus |
| • Dorsal Longitudinal Fasciculus | • Inferior Longitudinal Fasciculus |
| • Ventral Longitudinal Fasciculus | • Corpus Callosum |
| • Medial Lemniscus | • Anterior Commissure |
| • Central Tegmental Tract | • Fornix |
| • Cerebellothalamic Tract | • Inferior Fronto-Occipital Fasciculus |
| • Fasciculus Cinguli (Cingulum) | • Fronto-occipital Fasciculus |
| • Optic Radiation | • Superior Occipitofrontal Fasciculus |
| • Fasciculus Solitarius (Solitary Tract) | • Mammillothalamic Tract |
| • Medial Forebrain Bundle | • Anterior Thalamic Radiation |
| • Reticulospinal Tract | • Perforant Path |
| • Vestibulospinal Tract | • Commissural Fibers of the Hippocampus |
| • Lateral Spinothalamic Tract | • Thalamic Radiations |
| • Ventrolateral Spinothalamic Tract | • Corticopontine Tract |
| • Ventral Trigeminothalamic Tract | |
| • Dorsal Trigeminothalamic | |





From Chaos ...Comes Order

Unclear nomenclature for Association Pathways

Proposal of seven white matter *systems* defined by anatomical corridors

- Superior longitudinal system (SLS)
- Inferior longitudinal system (ILS)
- Middle longitudinal system (MidLS)
- Basal longitudinal system (BLS)
- Mesial longitudinal system (MesLS)
- Anterior transverse system (ATS)
- Posterior transverse system (PTS)





◦ Longitudinal system

◦ Superior (SLS)

◦ Inferior (ILS)

◦ Middle (MidLS)

◦ Basal (BLS)

◦ Mesial (MesLS)

◦ Transverse system

◦ Anterior (ATS)

◦ Posterior (PTS)

Spatial Awareness/Attention

Object Awareness/Attention

Comprehensive Sound

Comprehensive Vision

Comprehensive eMotivation

Articulated Language

Visual Language

ATS
PTS



SLS


i.e. Superior Longitudinal Fasciculus (SLF) and Arcuate Fasciculus (AF)

Connects frontal lobe to parietal, occipital and temporal lobes

- Travels with corona radiata above sulcus of the insula

Can be divided into four subsystems

- SLS-I i.e. superior SLF
- SLS-II i.e. middle SLF
- SLS-III i.e. inferior SLF
- SLS-IV i.e. AF



SLS I-III

i.e. Superior Longitudinal Fasciculus


Connecting the frontal lobe to the parietal lobe

- SLF I - Connect *superior* frontal to *superior* parietal lobe
- SLF II - Connect *middle* frontal with *middle* parietal lobe
 - angular [BA39] and *posterior* supramarginal gyrus [BA40]
- SLF III - Connect *inferior* frontal and *middle* parietal lobe
 - *anterior* supramarginal gyrus [BA40]

Dorsal Stream of Information i.e. WHERE?


Dominant – Language awareness/attention

Non-Dominant – Visuospatial awareness/attention




SLS I-III

Dorsal Stream – WHERE?



Visuospatial Circuit – *Where to look at?*






Language Circuit – *Where to move my mouth/conversation?*

SLS I-III

Dorsal Stream – WHERE?

51-year-old s/p hemorrhage right putamen with IVH s/p EVD and craniectomy developed Balint syndrome 2/2 b/l SLP dysfunction



60-year-old s/p left fronto-parietal lobe meningioma resection with IVH developed ideomotor apraxia

SLS IV

i.e. Arcuate Fasciculus

Connecting the inferior frontal to the superior temporal lobe

- i.e. Wernicke's connects with Broca

Direct/Long Segment = Repetition



- Sensory-motor mapping

Anterior Segment - Fluency

- Lexical-Semantic Processing - Basic comprehension ie words in context

Posterior Segment = Comprehension

- Lexical-Semantic Retrieval - Complex comprehension ie sentence



SLS IV

i.e. Arcuate Fasciculus

Neurodivergent structural anomalies in the left arcuate fasciculus in adults with developmental dyslexia

- left arcuate fasciculus = grapheme-phoneme decoding
- left inferior fronto-occipital fasciculus = direct word access

> [Brain](#). 2012 Mar;135(Pt 3):935-48. doi: 10.1093/brain/awr363. Epub 2012 Feb 10.

A tractography study in dyslexia: neuroanatomic correlates of orthographic, phonological and speech processing



Maaïke Vandermosten ¹, Bart Boets, Hanne Poelmans, Stefan Sunaert, Jan Wouters, Pol Ghesquière

Affiliations + expand


PMID: 22327793 DOI: [10.1093/brain/awr363](#)

RESEARCH ARTICLE

Open Access

Relation between aphasia and arcuate fasciculus in chronic stroke patients


Hyung Jun Tak and Sung Ho Jang*



Dominant AF most commonly effected in conduction aphasia i.e. fluent paraphasic repetition/spelling/writing impaired.

SLS I-IV


Non-Fluent Aphasia Agrammatic > Logopenic




BRAIN
A JOURNAL OF NEUROLOGY

White matter damage in primary progressive aphasias: a diffusion tensor tractography study

Sebastiano Galantucci,^{1,2} Maria Carmela Tartaglia,¹ Stephen M. Wilson,^{1,3} Maya L. Henry,¹ Massimo Filippi,² Federica Agosta,² Nina F. Dronkers,^{4,5} Roland G. Henry,⁶ Jennifer M. Ogar,¹ Bruce L. Miller¹ and Maria Luisa Gorno-Tempini^{1,7}





- SLF II no significant asymmetry
 - SLF II more dorsal>ventral connectivity in PFC
- SLF III **Non-Dominant** lateralization
 - SLF III more ventral>dorsal connectivity in PFC
 - improved fine control of complex actions through space and time? Social cognition?
- Arcuate Fasciculus **Dominant** Lateralization
 - improved fine sensory-speech mimicry

Virtual dissection and comparative connectivity of the superior longitudinal fasciculus in chimpanzees and humans

[Erin E. Hecht](#),^{1,2} [David A. Gutman](#),³ [Bruce A. Bradley](#),⁴ [Todd M. Preuss](#),⁵ and [Dietrich Stout](#)⁶

ILS


i.e. uncinate fasciculus (UC) and Inferior Fronto-occipital fasciculus (IFOF)

connects temporal/parietal/occipital lobes to frontal lobe

- pass below inferior limiting **sulcus** of the insula

Can be divided into four subsys

- ILS-I i.e. IFOF
- ILS-II i.e. IFOF
- ILS-III i.e. IFOF
- ILS-IV i.e. UC



ILS I-III

i.e. Inferior Fronto-occipital fasciculus (IFOF)

connects temporal/parietal/occipital lobes to frontal lobe



- ILS I – terminates medial PFC
- ILS II – terminates middle PFC
- ILS III – terminates lateral PFC

Ventral Stream of Information – WHAT?

ILS I dlPFC/pre-FOF = visual switching tasks




ILS II – vmPFC = goal-oriented behavior

ILS III – vlPFC = semantic processing i.e. object value



ILS I-III

Ventral Stream – WHAT?




Visuospatial Circuit – *What to look at?*

Language Circuit – *What am I looking at?*

ILS I-IV

Semantic Dementia/Lewy Body Dementia



JOURNAL ARTICLE

Language networks in semantic dementia

Federica Agosta, Roland G. Henry, Raffaella Migliaccio, John Neuhaus, Bruce L. Miller, Nina F. Dronkers, Simona M. Brambati, Massimo Filippi, Jennifer M. Ogar, Stephen M. Wilson ... [Show more](#)

Brain, Volume 133, Issue 1, January 2010, Pages 286–299,



<https://doi.org/10.1093/brain/awp233>

Published: 16 September 2009 [Article history](#)

BRAIN
A JOURNAL OF NEUROLOGY

White matter damage in primary progressive aphasia: a diffusion tensor tractography study

Sebastiano Galantucci,^{1,2} Maria Carmela Tartaglia,¹ Stephen M. Wilson,^{1,3} Maya L. Henry,¹ Massimo Filippi,² Federica Agosta,² Nina F. Dronkers,^{4,5} Roland G. Henry,⁶ Jennifer M. Ogar,¹ Bruce L. Miller¹ and Maria Luisa Gorno-Tempini^{1,7}




ILS IV

i.e. Uncinate Fasciculus (UC)

connects orbitofrontal/inferior frontal
with the anterior temporal lobe

Mnemonic associations (e.g. name +
face + voice = human) to modify
behavior/biasing of decisions

- Dominant - Language processing
- Non-Dominant Emotional understanding and empathy



ILS IV


Epilepsy/bvFTD Dementia

Uncinate seizures/Uncinate Fits

- Olfactory/Gustatory Hallucinations
- “Dreaming state”
- Oropharyngeal Automatism

FTLD; behavior and semantic subtypes

- Loss of empathy/sympathy/social inter-relatedness
- Oropharyngeal fixation



BLS


i.e. Inferior longitudinal fasciculus (ILF)

connects ventral temporal and occipital lobes

- pass below the inferior limiting sulcus of the insula
- consists of successive series of U-fibers

Can be divided into four subsystems

- BLS I - temporal cortex to lateral occipital (lateral)
- BLS II - temporal cortex to cuneus (medial)
- BLS III - temporal cortex to lingual gyrus (medioventral)
- BLS IV - temporal cortex to fusiform gyrus (ventral)



BLS



i.e. Inferior longitudinal fasciculus (ILF)

Hierarchy of Visuospatial Object Processing/ Comprehension

- Basic visual processing via BLS II
- Distance and depth perception, color determination via BLS I
- Letters and logical conditions (i.e., logical order) via *BLS III*
- face perception, object recognition, and reading via BLS IV

Clinical Implications

- Parkinson Disease/Lewy Body Disease Dementia
- Posterior Cortical Atrophy/Anton Syndrome



Visuospatial Ventral Stream
What is this?

BLS

Lewy Body Dementia

[Brain Behav.](#) 2020 Dec; 10(12): e01883.

Published online 2020 Oct 20. doi: [10.1002/brb3.1883](#)



PMCID: PMC7749587

PMID: [33078912](#)

Visual hallucinations and inferior longitudinal fasciculus in Parkinson's disease

[Natsuko Yuki](#),^{1,2} [Akira Yoshioka](#),^{2,3} [Ryo Mizuhara](#),¹ and [Tadashi Kimura](#)¹

► [Author information](#) ► [Article notes](#) ► [Copyright and License information](#) ► [PMC Disclaimer](#)



MesLS


i.e. Cingulum

connects prefrontal cortex and cingulate gyrus to hippocampus

- medial surface of the hemisphere
- arching from the frontal pole up to the amygdala area.

Can be divided into two subsystems

- MesLS I or “inner cingulum”
- MesLS II or “outer cingulum” or supracingulate



MesLS

i.e. Cingulum


Multimodal self-referential data relay

- **Anterior**; emotional response regulation, reward expectation, error monitoring
- **Posterior**: memory retrieval and spatial navigation
- **Lateral**: attention, decision making, language processing, and number processing


Clinical Significance

- Dementia NOS; AD, FTLD, PD
- Psych NOS;
ADHD/Schizophrenia/PTSD/OCD/ASD/BD

So it effects everything?



Most commonly defined as resting state data involved in self-awareness, introspection, and planning



MesLS

i.e. Cingulum

[Brain](#). 2016 Dec; 139(12): 3137–3150.

Published online 2016 Oct 22. doi: [10.1093/brain/aww241](https://doi.org/10.1093/brain/aww241)

PMCID: PMC5382939

PMID: [27797805](https://pubmed.ncbi.nlm.nih.gov/27797805/)


Disconnection between the default mode network and medial temporal lobes in post-traumatic amnesia

[Sara De Simoni](#),^{aww241-1} [Patrick J. Grover](#),^{aww241-1} [Peter O. Jenkins](#),^{aww241-1} [Lesley Honeyfield](#),^{aww241-2}
[Rebecca A. Quest](#),^{aww241-2} [Ewan Ross](#),^{aww241-1} [Gregory Scott](#),^{aww241-1} [Mark H. Wilson](#),^{aww241-3} [Paulina Majewska](#),^{aww241-1}
¹ [Adam D. Waldman](#),^{aww241-2} [Maneesh C. Patel](#),^{aww241-2} and [David J. Sharp](#)^{aww241-1}

► [Author information](#) ► [Article notes](#) ► [Copyright and License information](#) [PMC Disclaimer](#)

Post-traumatic amnesia (PTA) and traumatic brain injury (TBI) seem to be linked to transient/chronic disruptions in Default Mode Network

- connectivity with the parahippocampus
- connectivity between posterior (PCC, precuneus, lateral parietal) cortex and ventromedial prefrontal cortex (vmPFC).



MidLS

i.e. middle longitudinal fasciculus


Connects anterior temporal to inferior parietal lobe/Extrastriate area

- Posterior-dorsal segment; i.e. posterior STG
- Antero-ventral segment; i.e. anterior STG

Function

- Inconsistent evidence
- Attention biases?
- Motion detection?
- Sublexical representations?
- Repetition-phonological network?
- Naming?

Clinical Significance unclear but implicated in PPAs



ATS

i.e. Frontal Aslant Tract

Connects the superior to inferior prefrontal lobes



- Medial/superior frontal cortex (SMA/preSMA)
- Inferior frontal gyrus (Broca; pars triangularis/opercularis)

Can be divided into four subsystems

- ATS I - frontal area to anterior pars triangularis (Brodmann 47)
- ATS II - frontal area to posterior pars triangularis (Brodmann 45)
- ATS III - pre-SMA to pars opercularis (Brodmann 44)
- ATS IV - SMA to precentral gyrus

Function

- Language; expressive aphasia/speech apraxia/syntax



ATS

i.e. Frontal Aslant Tract

Functions

- Language order and syntax via ATS I
- translate from secondary or tertiary language via ATS II
- Inflectional morpheme via ATS III
- Nociception via ATS III
- Articulation via ATS IV

Clinical Implications


- Expressive Aphasia
- nfPPA/lvPPA



TactusTherapy

PTS

i.e. vertical temporo-parietal and occipital fasciculus




connects parietal occipital and posterior temporo-occipital


Can be divided into two subsystems

- PTS I i.e. temporo-parietal aslant tract
 - middle temporal gyrus to supramarginal/angular
- PTS II i.e. vertical occipital fasciculus
 - ventral transverse/temporo-occipital to intraparietal sulcus

Primary link between the dual visual processing streams

- VOF – Reading?
 - Pure word blindness, also known as pure alexia
- VTPF – visual word and language comprehension?






- Spatial Awareness/Attention*
- Object Awareness/Attention*
- Comprehensive Sound*
- Comprehensive Vision*
- Comprehensive eMotivation*

- Articulated Language*
- Visual Language*

Next time you see



Don't roll your eyes

Think

Less Focal-Symptomatology

And

More Disconnection-Phenomenology



Special Thanks to the
Ochsner Neurocognitive Team



References

