



Cervical Transforaminal Injections

What is transforaminal injection?

Transforaminal injection can help relieve pain in your neck and arm pain caused by herniated discs or other neck problems.

You have nerves that run from your spinal cord out to your neck, shoulder and arms. When one of those nerve “roots” (the end closest to your spinal cord) gets irritated and inflamed due to a damaged disc or other condition, it causes pain that can radiate from your neck to the arm and sometimes all the way to the hand.

A steroid medication can help reduce the inflammation and alleviate the pain. It’s injected into your neck, into the area between your spine and spinal cord.

How is transforaminal injection done?

First, you may be given an intravenous medication to relax you. Then, you’ll lie tilted on your side on an x-ray table.

The doctor will numb an area of skin on the side of your neck with a local anesthetic. Then, guided by an x-ray, he or she will:

- Insert a thin needle into your neck, targeting the specific nerve root causing your pain
- Inject a mixture of anesthetic (for temporary pain relief) and steroid (for longer term relief)

Usually, the procedure takes less than 5 minutes. Preparing for it will require your presence for at least one hour at the facility. You can go home the same day.

How effective is transforaminal injection?

Some patients report pain relief within 30 minutes after the injection, but pain may return a few hours later as the anesthetic wears off. Longer term relief usually begins in two to three days, once the steroid begins to work.

How long the pain relief lasts is different for each patient. For some, the relief lasts several months or longer. If the treatment works for you, you can have periodic injections to stay pain-free.

What are the risks?

The risk of complication from a transforaminal injection is very low. Rarely, complications include bleeding or infection at the injection site.

Side effects can include but not limited to:

- Numb or weak
- Dizziness
- Headache

These effects resolve within a few hours.

What happens after the procedure?

You can continue your regular diet and medications immediately. Do not drive or do any rigorous activity for 24 hours. Take it easy. You can return to your normal activities the next day.

It may take up to a week for the steroid to begin working. If you don't feel better within 14 days, see your doctor for more evaluation and to discuss different treatment.

Is transforaminal injection right for you?

Transforaminal injection may be right for you if your neck and arm pain has lasted longer than four weeks, is severe or has not improved with other treatments, such as physical therapy.

Talk to your physician about your symptoms and he/she will help decide what is right for you.